

2023 INTERNATIONAL CONGRESS OF ACTUARIES



BRIDGE TO TOMORROW

28 MAY – 1 JUNE 2023 • SYDNEY



**Actuaries
Institute
Australia**



**International Actuarial Association
Association Actuarielle Internationale**



Mortality analysis of mental health cohorts with co-morbid physical conditions: A case study from Australia

Maggie Lee

© The Australian National University, Commonwealth Department of Health and Aged Care

*This presentation has been prepared for the 2023 International Congress of Actuaries.
The Actuaries Institute Australia Council wishes it to be understood that opinions put forward herein are not necessarily those
of the Institute and the Council is not responsible for those opinions.*



Outline

1. Introduction
2. Data and methodology
3. Results
4. Summary



Background

- Mental Health: a key priority for many governments around the world
- Increasing **prevalence**
 - 13% increase in last decade to 2017 (WHO, 2022)
- Increasing **social and health consequences**
 - Cost to the world economy: US\$2.5 trillion in 2010 increasing to US\$6 trillion by 2030 (Health T.L.G, 2020)
 - Reduction in life expectancy of 1.4 to 32 years, with median of 10.1 years (Walker et al. 2015).
- Those with mental health conditions also tend to have **physical comorbidities**
 - **Chronic conditions:** cardiovascular and respiratory diseases, cancers, diabetes
 - Number of life years lost due to physical conditions for individuals with mental health conditions is increasing (Larsen et al. 2018, Erlangsen et al. 2017)



Setting context – Policy relevance

Improving the physical health and life expectancy for people living with mental health conditions is a focus area for both the:

- National Mental Health and Suicide Prevention Agreement and
- Fifth National Mental Health and Suicide Prevention Plan (Priority 5)

The research under HMHB (DoHAC) will contribute to the evidence base to inform activities that are required under these government arrangements.



Life expectancy gaps and preventable risks

- Work to date has focussed on HMHB objective 1
 - What are the **health and social outcomes** of individuals that have co-morbid mental and physical health issues (for different diagnosis groups, compared to those without conditions)?
 - To get a better understanding of the health and social outcomes, the aim of study is threefold:
 1. **Quantify the life expectancy gap** of individuals in Australia that have mental health conditions and the factors including physical comorbidities that contribute to premature mortality
 2. Identify the risk factors for mental health cohorts compared to those without conditions. Examples of factors investigated include smoking, diet, physical activity and alcohol consumption.
 3. Identify preventable hospitalisations and hospital re-admissions for both mental and physical health issues



Outline

1. Introduction
- 2. Data and methodology**
3. Results
4. Summary




Data

- Australian Institute of Health and Welfare's (AIHW) National Integrated Health Services Information Analysis Asset (NIHSI AA)
 - Data included from July 2010 to June 2017
 - [Hospitalisation](#), Medicare Benefits Schedule (MBS), Pharmaceutical Benefits Scheme (PBS), Residential Aged Care, National Death Index
- Australian Bureau of Statistics' (ABS) Multi-Agency Data Integration Project (MADIP)
 - Data included from January 2011 to December 2020
 - MBS, PBS, National Death Index, [Census/Demographics, Migration, Survey \(National Health Survey and Survey of Disability, Ageing and Carers\)](#) and [Income/Taxation](#) data



Methods

- Design: Population-based cohort study (Broad Mental Health and SMI)
- Mortality Rate Ratios (MRR): Poisson model adjusting for sex, age
- Life expectancy: Period life table method
- Life Years Lost (LYL): Differences in life expectancy after disease onset
 - Only for SMI



Source: Plana-Ripoll (2022)

- Weighted average excess Life-Years Lost: weighted average life years lost using age-of-onset distribution
- Health Expectancy: Sullivan method



Outline

1. Introduction
2. Data and methodology
- 3. Results**
4. Summary



Life expectancy gaps – Exploratory analysis

	No. of people	Person-years	Deaths	Death rate	Reference*	Mortality rate ratio
Broad mental health cohort	7,436,074	15,501,773	273,861	17.7	10.8	1.6
Female	4,339,742	9,222,520	143,389	15.5	10.3	1.5
Male	3,096,332	6,279,253	130,472	20.8	10.8	1.9
	No. of people	Person-years	Deaths	Death rate	Reference*	Mortality rate ratio
SMI cohort	846,210	1,454,831	82,466	56.7	14.6	3.9
Female	426,133	731,866	43,446	59.4	16.5	3.6
Male	420,077	722,965	39,020	54.0	12.3	4.4


*Reference is the standardised death rate of the general Australian population to the age and gender distribution of the broad mental health and SMI cohorts



Life expectancy gaps – Exploratory analysis

- Mortality rate of broad mental health and SMI cohorts compared the general Australian population by gender

Graph: Mortality rate of broad mental health and SMI cohorts compared to the general Australian population (NIHSI AA)





Life expectancy gaps – Preliminary results (Life Table)


Life expectancy of the general Australian population to the broad mental health cohort and SMI cohort

	Male						Female					
	0	15	45	65	85+		0	15	45	65	85+	
Age												
All Australians*	80.4	65.8	37.1	19.5	4.7		84.5	69.9	40.6	22.3	5.3	
Broad mental health cohort	70.0	56.4	29.2	13.6	0.5		75.3	62.4	33.8	16.4	0.5	
SMI cohort	58.8	45.8	21.0	8.3	0.5		63.7	50.6	24.4	10.6	0.5	
Life expectancy gap for Broad Mental Health cohort	10.4	9.4	7.9	5.9	4.2		9.2	7.5	6.8	5.9	4.8	
Life expectancy gap for SMI cohort	21.6	20	16.1	11.2	4.2		20.8	19.3	16.2	11.7	4.8	



Life expectancy gaps – Preliminary results (Sullivan method)

Graph: Proportion of remaining life spent in disease state at each age (NIHS/AA)






- Broad mental health
 - Males: 19% to 31%
 - Females: 27% to 44%
- SMI
 - Males: 0% to 5%
 - Females: 0% to 7%




Life expectancy gaps – Preliminary results

(Life-years lost)



**All Australian is using NHI data so won't match to ABS lifetable at this stage*



Onset distribution	Total average Excess Life-years lost	Internal causes	External Causes
New Cases – FY14FY17	11.7	9.8	1.9
New Cases – FY15FY17	11.6	9.7	1.9
New Cases – FY16FY17	11.6	9.7	1.9
New Cases – FY17	11.6	9.7	1.9

Outline

1. Introduction
2. Data and methodology
3. Results
- 4. Summary**



Summary

- **Mortality rate ratio**
 - 1.6 times higher for broad mental health and 3.9 times higher for SMI
 - Broadly consistent with international studies
 - » Plana-Ripoll et al. (2019): MRR range from 1.9 to 3.1 across all genders in Denmark
 - » Pan et al. (2020): MRR range from 1.6 for depression, 2.4 for bipolar and 3.7 for schizophrenia across all genders in Taiwan
 - » Krupchanka et al. (2018): MRR range from 1.6 (mood disorders), 2.3 (schizophrenia) and 3.5 (substance use disorders) in Czech Republic
 - » Ajetunmobi et al. (2013): MRR range from 1.5 to 4.4 across all genders in Scotland
- **Life expectancy gap**
 - At age 15 (Broad Mental Health: 9.4 years – Males/7.5 years – Females; SMI: 20 years – Males/19.3 – Females)
 - » Lawrence et al. (2013): 15.9 years for males and 12 years for females in 2005 for psychiatric patients in W.A.
 - » Nordentoft et al. (2013): 20 years for males and 15 years for females from 2000 to 2006 for patients with psychiatrist disorder
- **Excess life years lost** – weighted average was 11.6 years
 - » Erlangsen et al. (2017): 10.2 years for males and 7.3 years for females across all mental disorders in Denmark
 - » Plana-Ripoll et al. (2019): 12.1/9.4 for schizophrenia, 7.9/6.2 for mood disorders and 8.6/6.6 for neurotic disorders in Denmark



Further research

- Further segmentation of mental health cohorts by different diagnosis groups and by gender
- Exploration on best methods to ensure the most accurate and practical excess mortality measures have been explored
- Further analysis into the association between mental health conditions and physical health conditions
- Exploration of risk stratification tools (similar to that developed by Khanna et al. 2019) to help inform preventable hospitalisations and hospital re-admissions

BRIDGE TO TOMORROW
2023 INTERNATIONAL
CONGRESS OF ACTUARIES
28 MAY – 1 JUNE 2023 · SYDNEY



Thank you



Appendix – Broad mental health cohort definition

- Broad mental health cohort definition (based on AIHW)
 - **Hospital** primary diagnosis using ICD-10 Codes: F00 – F99
 - » This includes F10-F19 (substance use disorders), F20-F29 (Schizophrenia), F30-F39 (Mood disorders), F40-F48 (Neurotic disorders), F50 (Eating disorders), F60 (Personality disorders), F70-F79 (Intellectual disabilities), F84 (Development disorders), F90-F98 (Behavioural disorders).
 - **MBS items** related to mental health services
 - » Includes services provided psychiatrists (initial consultation, patient attendance, group psychotherapy, COVID-19 Services etc); GPS (mental health treatment plan, Eating disorders treatment plan, COVID-19 mental health services, focussed psychological strategies); Clinical psychologists (psychological treatment services, eating disorders psychological treatment, focussed psychological strategies, COVID-19); Psychologists including clinical psychologists (Enhanced primary care, Allied health services for Indigenous Australians); Other Allied Health providers (Enhanced primary care) etc.
 - **PBS items** (ATC codes) related to mental health services
 - » antipsychotics (N05A), anxiolytics (N05B), hypnotics and sedatives (N05C), antidepressants (N06A), and psychostimulants, agents used for ADHD and nootropics (N06B)— prescribed by all medical practitioners (GPs, non-psychiatrist specialists and psychiatrists).



Appendix – SMI cohort definition

- The SMI cohort has been defined as any individual who meets at least one of the following conditions for each financial year:
 1. They were prescribed anti-psychotic medication. Anti-psychotic medication includes N05A excluding N05AB04; or
 2. They were prescribed anticonvulsant/mood stabilising medication and they saw a psychologist or clinical psychologist more than 4 times. Anticonvulsant/mood stabilising medication includes sodium valproate N03AG01 or carbamazepine N03AF01 or lamotrigine N03AX09); or
 3. They saw a psychiatrist more than 4 times; or
 4. Were hospitalised with an ICD-10-AM primary diagnosis of Schizophrenia/Schizoaffective disorder (F20, F25 ICD-10-AM codes) or bipolar disorder (F31 ICD-10-AM code)



Appendix – Possible age of onsets

Mental disorders Examples of diagnoses included in each group	Abbreviated name used in this study	ICD-10	ICD-8 equivalency	Earliest possible age at onset (years)
Any mental disorder	Any disorder	F00 – F99	290 – 315	1
Organic, including symptomatic, mental disorders Includes dementia in Alzheimer's disease, vascular dementia, etc.	Organic disorders	F00 – F09	290-09, 290-10, 290-11, 290-18, 290-19, 292 x9, 293 x9, 294 x9, 309 x9	35
Mental and behavioral disorders due to psychoactive substance use Includes use of alcohol, cannabis, cocaine, nicotine, opioids, sedatives, hypnotics, anxiolytics, etc.	Substance use disorders	F10 – F19	291 x9, 294-39, 303 x9, 303-20, 303-28, 303-90, 304 x9	10
Schizophrenia and related disorders Includes schizophrenia, schizotypal disorders, schizoaffective disorders and other psychotic disorders.	Schizophrenia	F20 – F29	295 x9, 296-89, 297 x9, 298-29, 298-99, 299-04, 299-05, 299-09, 301-83	10
Mood disorders Includes bipolar disorder, depressive disorders, etc.	Mood disorders	F30 – F39	296 x9 (excluding 296-89), 298-09, 298-19, 300-49, 301-19	10
Neurotic, stress-related, and somatoform disorders Includes anxiety disorders, phobias, obsessive-compulsive disorders, etc.	Neurotic disorders	F40 – F48	300 x9 (excluding 300-49), 305 x9, 305-68, 307-99	5
Eating disorders Includes anorexia nervosa, bulimia nervosa, etc.	Eating disorders	F50	305-60, 306-50, 306-58, 306-59	1



Appendix – Possible age of onsets

Eating disorders <i>Includes anorexia nervosa, bulimia nervosa, etc.</i>	Eating disorders	F50	305·60, 306·50, 306·58, 306·59	1
Specific personality disorders	Personality disorders	F60	301·x9 (excluding 301·19), 301·80, 301·81, 301·82, 301·84	10
Intellectual disabilities	Intellectual disabilities	F70 – F79	311·xx, 312·xx, 313·xx, 314·xx, 315·xx	1
Pervasive developmental disorders <i>Includes autism spectrum disorder</i>	Developmental disorders	F84	299·00, 299·01, 299·02, 299·03	1
Behavioral and emotional disorders with onset usually occurring in childhood and adolescence <i>Includes attention-deficit hyperactivity disorder, conduct disorders, childhood emotional disorders, etc.</i>	Behavioral disorders	F90 – F98	306·x9, 308·0x	1

Source: Plana-Ripoll et al. (2019)